

## Footnotes for the Chart: Soluble Fiber Foods

Chart available at:

[www.gardeningplaces.com/articles/](http://www.gardeningplaces.com/articles/)

[1] Li et al., Individual Sugars, Soluble, and Insoluble Dietary Fiber Contents of 70 High Consumption Foods; *Journal of Food Composition and Analysis*. Volume 15, Issue 6, December 2002, Pages 715–723, Table 1.

<http://www.sciencedirect.com/science/article/pii/S0889157502910963>

[2] Manzi et al., Nutritional value of mushrooms widely consumed in Italy; *Food Chemistry*. Volume 73, Issue 3, May 2001, Pages 321–325.

<http://www.sciencedirect.com/science/article/pii/S0308814600003046>

[3] Ramulu and Rao, Total, insoluble and soluble dietary fiber contents of Indian fruits; *Journal of Food Composition and Analysis*. Volume 16, Issue 6, December 2003, Pages 677–685.

<http://www.sciencedirect.com/science/article/pii/S0889157503000954>

[4] Reyes-Caudillo et al., Dietary fibre content and antioxidant activity of phenolic compounds present in Mexican chia (*Salvia hispanica* L.) seeds; *Food Chemistry*. Volume 107, Issue 2, 15 March 2008, Pages 656–663.

<http://www.sciencedirect.com/science/article/pii/S0308814607008709>

[5] Tosi et al., Dietary fiber obtained from amaranth (*Amaranthus cruentus*) grain by differential milling; *Food Chemistry*. Volume 73, Issue 4, June 2001, Pages 441–443.

<http://www.sciencedirect.com/science/article/pii/S0308814600003265>

[6] Repo-Carrasco-Valencia et al., Dietary fiber and other functional components in two varieties of crude and extruded kiwicha (*Amaranthus caudatus*); *Journal of Cereal Science*. Volume 49, Issue 2, March 2009, Pages 219–224.

<http://www.sciencedirect.com/science/article/pii/S0733521008001689>

[7] Repo-Carrasco-Valencia and Serna, Quinoa (*Chenopodium quinoa*, Willd.) as a source of dietary fiber and other functional components; *Food Science and Technology (Campinas)*. *Ciênc. Tecnol. Aliment.* vol.31 no.1 Campinas Jan./Mar. 2011.

[http://www.scielo.br/scielo.php?pid=S0101-20612011000100035&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S0101-20612011000100035&script=sci_arttext)

[8] Repo-Carrasco-Valencia et al., Chemical and Functional Characterization of Kañiwa (*Chenopodium pallidicaule*) Grain, Extrudate and Bran; *Plant Foods for Human Nutrition*. June 2009, Volume 64, Issue 2, p. 94-101.

<http://link.springer.com/article/10.1007/s11130-009-0109-0>

[9] Steadman et al., Buckwheat Seed Milling Fractions: Description, Macronutrient Composition and Dietary Fibre; *Journal of Cereal Science*. Volume 33, Issue 3, May 2001, Pages 271–278.

<http://www.sciencedirect.com/science/article/pii/S0733521001903662>

- [10] Todd Dalotto, *The Hemp Cookbook: From Seed to Shining Seed* (Rochester, VT; Healing Arts Press, 2000) p. 16; data provided by Don Wirtshafter.
- [11] Udo Erasmus, *'Fats that Heal, Fats that Kill'*, (Burnaby, BC, Canada: Alive Books, 1993), p. 289.
- [12] nutiva.com, "Organic Hemp Protein", retrieved July 2014;  
<https://store.nutiva.com/hemp-protein/>
- [13] Values for soluble and insoluble fiber based on 1:9 ratio and total fiber of 6% (per multiple sources).
- [14] Leila Picolli da Silva and Maria de Lourdes Santorio Ciocca, Total, insoluble and soluble dietary fiber values measured by enzymatic–gravimetric method in cereal grains; *Journal of Food Composition and Analysis*. Volume 18, Issue 1, February 2005, Pages 113–120. Note: values are dry matter average of several cultivars, adjusted by the author to a moisture content of 9%.  
<http://www.sciencedirect.com/science/article/pii/S0889157503001807>
- [15] Costa et al., Chemical composition, dietary fibre and resistant starch contents of raw and cooked pea, common bean, chickpea and lentil legumes; *Food Chemistry*. Volume 94, Issue 3, February 2006, Pages 327–330.  
<http://www.sciencedirect.com/science/article/pii/S0308814604008519>
- [16] S. Shobana and N.G. Malleshi, Preparation and functional properties of decorticated finger millet (*Eleusine coracana*); *Journal of Food Engineering*. Volume 79, Issue 2, March 2007, Pages 529–538.  
<http://www.sciencedirect.com/science/article/pii/S0260877406001877>
- [17] Ann L. Gerhardt and Noreen B. Gallo, Full-Fat Rice Bran and Oat Bran Similarly Reduce Hypercholesterolemia in Humans; *Journal of Nutrition*. May 1, 1998 vol. 128 no. 5 865-869.  
<http://jn.nutrition.org/content/128/5/865.full>
- [18] Sudha et al., Influence of fibre from different cereals on the rheological characteristics of wheat flour dough and on biscuit quality; *Food Chemistry*. Volume 100, Issue 4, 2007, Pages 1365–1370.  
<http://www.sciencedirect.com/science/article/pii/S0308814605010575>
- [19] Bunzel et al., Cell wall hydroxycinnamates in wild rice (*Zizania aquatica* L.) insoluble dietary fibre; *European Food Research and Technology*. June 2002, Volume 214, Issue 6, p. 482-488.  
<http://link.springer.com/article/10.1007/s00217-002-0512-3>
- [20] Bonafaccia et al., Characteristics of spelt wheat products and nutritional value of spelt wheat-based bread; *Food Chemistry*. Volume 68, Issue 4, March 2000, Pages 437–441.  
<http://www.sciencedirect.com/science/article/pii/S0308814699002150>
- [21] Shih et al., Effects of drying and extrusion on colour, chemical composition, antioxidant activities and mitogenic response of spleen lymphocytes of sweet potatoes; *Food Chemistry*. Volume 117, Issue 1, 1 November 2009, Pages 114–121. Note: values adjusted from dry matter weight to 3.0% total fiber in raw sweet potato (per USDA National Nutrient Database for Standard Reference, Release 26).  
<http://www.sciencedirect.com/science/article/pii/S0308814609004075>

[22] A. Redondo-Cuenca et al., Chemical composition and dietary fibre of yellow and green commercial soybeans (*Glycine max*); *Food Chemistry*. Volume 101, Issue 3, 2007, Pages 1216–1222. Note: values cited are for conventional commercial soybeans.

<http://www.sciencedirect.com/science/article/pii/S0308814606002445>

[23] Huang et al., Nutrient Composition of Taro Corms and Breadfruit; *Journal of Food Composition and Analysis*. Volume 13, Issue 5, October 2000, Pages 859–864.

<http://www.sciencedirect.com/science/article/pii/S0889157500909360>

[24] Deepa et al., Nutrient composition and physicochemical properties of Indian medicinal rice – Njavara; *Food Chemistry*. Volume 106, Issue 1, 1 January 2008, Pages 165–171.

<http://www.sciencedirect.com/science/article/pii/S0308814607005225>

[25] Susan Sungsoo Cho and Mark Dreher, editors, *Handbook of Dietary Fiber* (Marcel Dekker, Inc.: New York, 2001), Table 3.

[http://books.google.com/books?id=\\_XEQLhqtqxEC](http://books.google.com/books?id=_XEQLhqtqxEC)

Note from the book: "Based on published values: Am. Assoc. of Cereal Chemists, 1987; Dreher, 1987; Anderson and Bridges, 1988; Cardozo and Eitenmiller, 1988; Del Toma et al., 1988; Mongeau et al., 1989; USDA, 1989; Toma and Curtis, 1989; Ranhotra et al., 1990; Prosky and DeVries, 1992; and Marlett 1991."

[26] Takeshi Nagai and Toshio Nagashima, Functional Properties of Dioscorin, a Soluble Viscous Protein from Japanese Yam (*Dioscorea opposita* Thunb.) Tuber Mucilage Tororo; *Z. Naturforsch.* 61c, 792–798 (2006);

<http://znaturforsch.com/ac/v61c/s61c0792.pdf>